



Contents lists available at BioMedSciDirect Publications

## International Journal of Biological & Medical Research

Journal homepage: [www.biomedscidirect.com](http://www.biomedscidirect.com)



### Original Article

## The epidemiological features and laboratory diagnosis of keratomycosis

Anusuya devi. D, Ambica.R, Nagarathnamma T.

Department of Microbiology, Bangalore medical college and research institute, Bangalore -560002, Karnataka

#### ARTICLE INFO

##### Keywords:

Calcofluor white stain

Fusarium species

Potassium hydroxide mount

Mycotic keratitis

#### ABSTRACT

**Aims & objective:** To report the epidemiological features and laboratory diagnosis of 100 cases of fungal keratitis diagnosed at Regional Institute of Ophthalmology, Bangalore Medical College & Research Institute, Bangalore. **Methods:** A total of 100 patients with suspected fungal corneal ulcer satisfying the inclusion criteria were studied at Regional Institute of Ophthalmology, Bangalore Medical College & Research Institute, Bangalore during the period December 2009 to February 2011. Socio-demographic data, risk factors, seasonal variation and laboratory findings were studied. **Results:** Males (64) were affected significantly more than females (36). Of 100 patients, 78 were in the affected age group (20-50yrs). Ocular trauma was seen in 34 patients. There was a higher incidence of fungal keratitis during the monsoon and winter than summer. KOH, CFW and Gram stain revealed fungus in 46,51 and 17 patients respectively. Sensitivity of CFW was high (94.6%) followed by KOH (82.05%) and Gram stain (28.21%) respectively. Fusarium (42.1%), Aspergillus spp (28.93%) were predominantly isolated. **Conclusion:** keratomycosis is predominant in young adults with trauma as the major predisposing factor. Direct microscopy with CFW was more sensitive in detecting fungal elements. A simple and rapid Calcofluor White smear examination of corneal scraping early during the disease result in significant progress of the disease in initiating early and specific therapy.

© Copyright 2010 BioMedSciDirect Publications IJBMR -ISSN: 0976:6685. All rights reserved.

### 1. Introduction

Fungal keratitis is a suppurative, ulcerative and sight-threatening infection of the cornea that sometimes leads to loss of the eye. Corneal infection of fungal etiology is very common and represents 30%-40% of all cases of culture-positive infectious keratitis in South India.<sup>1</sup> WHO estimates global blindness about 20-40 million. In India there are approximately 10 million blind persons and 3 million are corneally blind<sup>2</sup>.

The majority of cases occur among agricultural workers following corneal trauma with vegetative matter contaminated by fungi. These are opportunistic organisms and colonize when natural defences of eye are abrogated by corneal trauma, use of topical steroids or any other predisposing factors.<sup>3</sup> Culture and direct microscopy are the two important investigations that are widely used. Culturing of microbial pathogens is considered to be the gold standard whereas direct microscopic evaluation of smears provides immediate information about the causative

organisms for initiating treatment.<sup>4,5</sup>

As the background knowledge of these factors accumulate, it is becoming important to identify the fungal corneal ulceration and to standardize the laboratory diagnostic methods so that an accurate early diagnosis can be made. Hence, this study aims to identify the most common causative organisms and also compares the efficacy of Calcofluor White staining with other methods like 10% KOH wet mount and Gram stain for early diagnosis of keratomycosis.

### 2. Objectives of the study

To evaluate the sensitivity, specificity and predictive value of Calcofluor White stain.

To compare the efficacy of Calcofluor White stain with 10% KOH wet mount and Gram stain.

To isolate and identify the fungal pathogens using 'c' streak method on Sabouraud's dextrose agar plate.

\* Corresponding Author : : **Dr. Anusuya devi.D,**  
Department of microbiology, Bangalore medical college and  
research institute, Bangalore -560002, Karnataka.  
E-mail- [annudocrajesh@rediffmail.com](mailto:annudocrajesh@rediffmail.com)  
Phone no: 9986088416.

### 3. Patients and methods

This study included 100 patients with clinically suspected fungal corneal ulcers attending the department of ophthalmology, Regional institute of Ophthalmology, Bangalore and the duration of study period was from December 2009 to February 2011. Inclusion Criteria:

1. Patients with corneal ulcer having loss of the corneal epithelium with underlying stromal infiltration and suppuration associated with signs of inflammation with or without hypopyon.
2. Patients who had received antimicrobial therapy and not responding to treatment were included in this study.

Patients with suspected (or) confirmed viral keratitis, bacterial keratitis, interstitial keratitis and sterile neurotropic ulcers were excluded.

#### Sample preparation

Multiple corneal scrapings were taken with the help of slit lamp using 26 gauge needle on a syringe. The corneal ulcer bed and margins were taken under topical Anaesthesia using 4% Lignocaine hydrochloride after removal of debris or discharge from the vicinity.<sup>6,7</sup>

#### Direct microscopic examination

The materials from the corneal scrapings were examined under 10% KOH, Calcofluor white stain and Gram stain.

- a. KOH preparation: wet mount preparation in 10% KOH was done to detect the presence of fungal elements. [Figure 1]
- b. Calcofluor white stain: one drop of calcofluor white stain (comprising 1g/L calcofluor white M2R and Evans blue Sigma-Aldrich, catalog No 18909 100 ml) and one drop of 10% KOH was added to the slide. The slide was then left to stand for 10 mts and was examined under fluorescence microscopy using blue light excitation (300-400 nm for the emission wavelength with excitation at around 355 nm).<sup>11</sup>[Figure 2]

- c. Gram stain: was done to look for hyphae, budding yeast cells and pseudohyphae.

#### Culture of specimens

The corneal scrapings obtained were inoculated directly onto Sabouraud's dextrose agar with antibiotics but without actidione. The material was directly inoculated onto the surface of Sabouraud's dextrose agar media in a row of c-shaped streaks and incubated aerobically at 25°C and 37°C. The plates were examined daily during the first week and twice weekly during the next two weeks.<sup>4,5</sup>The isolates were identified by standard laboratory procedures.<sup>4</sup>

The mycelial isolates were identified by their

Colony characteristic on SDA

Microscopic morphology on LPCB wet mount.

Slide culture

The yeast isolates were identified by standard tests

Germ tube test

Chlamyospore production on cornmeal agar

Growth on CHROMagar media.<sup>4</sup>

No growth was observed even after 3 weeks of incubation the culture was considered as sterile and the plates were discarded.<sup>4,8,9</sup>

#### Statistical methods

Chi-square /Fisher exact test has been used to find the significance of study. Statistical software: The Statistical software namely SAS 9.2, SPSS 15.0, Stata 10.1, MedCalc 9.0.1 ,Systat 12.0 and R environment ver.2.11.1 were used for the analysis of the data and Microsoft word and Excel have been used to generate graphs, tables etc.

**Table 1: Age and gender distribution**

Age group (years)	Male		Female		Total	
	No.	%	No.	%	No.	%
1-10	2	3.12	1	2.77	3	3
11-20	2	3.12	2	5.55	4	4
21-30	10	15.62	5	13.88	15	15
31-40	11	17.18	8	22.22	22	22
41-50	14	21.87	9	25.00	20	20
51-60	16	25.00	5	13.88	21	21
>60	9	14.06	6	16.66	15	15
Total	64	100	36	100	100	100%

**Table-2: Occupational distribution**

Occupation	No. of patients (%)
Farmers	44
Housewives	16
Industrial workers	15
Manual labourers	14
Business class	5
Unemployed	4
Students	2
Total	100

**Table- 3: Predisposing factor**

Predisposing factor	No of cases	Percentage (%)
Trauma	34	62.96
Topical steroids	14	25.92
Diabetes mellitus	1	1.85
Chronic dacryocystitis	1	1.85
Contact lenses	2	3.70
HIV	1	1.85
None	1	1.85
Total	54	100

**Table -4: Nature of trauma**

Nature of trauma	No of cases
Vegetative matter	34
Paddy	14
Thorn prick	1
Wooden sticks	1
Foreign body	2
a. Lime injury	1
b. Stone	1
c. Dust	54
d. Finger nail injury	
Total	

**Table-5: Direct demonstration of organisms**

Method	Positive	Percentage	Negative	Percentage
KOH	46	46%	54	54%
CFW	51	51%	49	49%
Gram stain	17	17%	83	83%

**Table- 6: Correlation of KOH, CFW, Gram with Culture:**

Method	Direct microscopy positive Culture Positive	Direct microscopy positive Culture negative	Direct microscopy negative Culture Positive	Direct microscopy negative Culture negative	Total
KOH	32	14	7	47	100
CFW	36	13	5	46	100
Gram stain	11	6	28	55	100


**Table -7: Comparison of sensitivity, specificity, positive predictive value and negative predictive values.**

Method	Sensitivity	Specificity	PPV	NPV	Accuracy	P value
KOH	82.05	77.05	69.57	87.04	79.00	<0.001**
CFW	94.06	75.41	69.38	90.20	80.00	<0.001**
GRAM	28.21	90.16	64.71	66.27	66.00	0.017*

**Table -8: Distribution of isolates in fungal keratitis**


Species	No of isolates	%
<i>Fusarium species</i>	16	42.10
<i>Aspergillus fumigatus</i>	06	15.78
<i>Aspergillus flavus</i>	03	7.89
<i>Aspergillus niger</i>	02	5.26
<i>Acremonium species</i>	02	5.26
<i>Curvularia</i>	02	5.26
<i>Cladosporium</i>	01	2.63
<i>Penicillium species</i>	03	7.89
<i>Candida albicans</i>	03	7.89
Total	38	100%

**Figure 1-KOH wet mount**



**Figure 1 showed fungal elements in KOH wet mount**

**Figure 2-CFW mount**



**Figure 2 showed fungal elements in CFW mount.**

#### 4. Results:

Of the 100 cases studied, 64(64%) were males and 36(36%) were females. Male to female ratio was 1.76:1. The age distribution showed the incidence of fungal keratitis predominantly between the 40- 60 yrs i.e. the working period of life. The mean age was 40.85 years [Table-1].

Fungal keratitis was predominantly seen in people engaged in farming (44%) followed by housewives (16%), industrial workers (15%) and labourers (14%) respectively. Agriculture is the most common occupation predisposing to fungal corneal ulcers [Table-2]. Out of the 100 patients, 68 belonged to low socio economic strata while 32 patients were from middle socio economic status. The majority of our patients were involved in agriculture and daily labourers. These persons were therefore exposed to a high risk of corneal trauma with vegetative matter while working in the fields.

Table 3 describes the occupational distribution of patients. In our study as many as 34 (62.96%) had corneal trauma as predisposing factor. The other risk factors were use of topical steroids (14), chronic dacryocystitis (1), HIV (1), Diabetes mellitus (1) and contact lens usage (2). As seen from the table 4, corneal trauma with vegetative matter seen in 20 (34.48%). Other causes of trauma were foreign body, stone, dust. Corneal trauma with vegetative matter was the most important risk factor for fungal keratitis.

Out of 100 cases studied, KOH was positive in 46 patients, CFW was positive in 51 patients and Gram stain positive in 17 patients. In the present study, positive fungal culture was taken as the gold standard showed the sensitivity of CFW (94.6) was higher in detection of fungal filaments compared to KOH (82.5) and Gram stain (28.21). Table 5,6,7 shows the fungal culture and direct microscopy methods and sensitivity & specificity of staining methods relative to fungal culture results.

Of the 38 cultures positive on SDA, *Fusarium* spp (42.1%) was the most common isolate followed by *Aspergillus* spp (28.93%) among which *Aspergillus fumigatus* was the predominant isolate. The other common isolates were *Curvularia* (5.26%), *Penicillium* spp. (7.89%), *Candida albicans* (7.89%), *Acremonium* spp, (5.26%) and *Cladosporium* spp (2.63%). The total percentage of culture positivity was 38%. *Curvularia* spp. (3.44%) was most common among the pigmented fungi followed by *Cladosporium* spp (1.72%) [Table-8].

#### 5. Discussion

Keratomycosis is an important ophthalmic problem in all parts of the world, as it leads to corneal blindness and sometimes loss of the vision. Its incidence is reported to vary from 7%- 40% in various parts of India.<sup>9,10</sup> Early and rapid identification of the pathological organism is the key to ensuring successful medical

therapy for corneal infections and is particularly true for fungal keratitis. Thus, microscopical evaluation of corneal scrapings from patients with suspected fungal keratitis is of the utmost importance because early and rapid detection of fungal structures permits early initiation of antifungal therapy<sup>11</sup>.

The mean age in the present study was noted to be 40.85yrs (20-50yrs). This is comparable with the study of Samar K Basak et al.<sup>5</sup> The age distribution showed the incidence of fungal keratitis predominantly between the 4th to 6th decades, reflecting the active working period of life and hence the increased vulnerability to injury during outdoor activities. The male predominance of fungal keratitis (64%) noted in the present study correlates with the studies of Bharathi et al, Srinivasan et al and Chowdhary et al.<sup>6,12,13</sup> Trauma was the major risk factor in 34% of the affected eyes.

The sensitivity of KOH used in the present study for diagnosing mycotic keratitis was determined to be (82.05%) and is comparable to the studies by Jagdish Chander et al and Tahereh Shokohi et al.<sup>14,15,16</sup> Calcoflour white was found to have a better sensitivity (94.6%) as compared to KOH and similar findings have been reported by Savithri Sharma et al, Tahereh Shokohi et al, Zhang et al and Usha Gopinathan et al.<sup>17,18,19</sup> Specificity of Gram stain, KOH and CFW stain were 90.16, 77.05 and 75.41% respectively.

Chander et al<sup>18</sup> in their study stated that CFW on direct microscopy detected fungi in 20 (95.2%) patients in comparison with 15 (71.4%) patients by both KOH wet mount and culture. They confirmed the superiority of KOH+CFW in comparison of with KOH alone and culture. Sharma et al<sup>19</sup> also confirmed the superiority of KOH+CFW stain in detection of fungal elements in corneal scraping. It is not only highly specific and sensitive but also has a high NPV and PPV for both early and advanced keratitis. Since they had seen some false positive results in advanced keratitis, they recommended resorting to antifungal therapy whenever a KOH+CFW stain smear is positive for fungus. Shokohi et al<sup>10</sup> in their study revealed a sensitivity of 71.4% for KOH and 42.9% for Gram stain. They concluded that KOH with CFW as the single most important screening test for rapid diagnosis of fungal corneal ulcer. Zhang et al<sup>11</sup> found out sensitivity of KOH+CFW was 96.6%. This combined staining method, enhancing the rate of correct diagnosis.

In the present study, 38 of the 100 corneal ulcer cases (38%) yielded fungal isolates on culture. Of these, *Fusarium* spp (42.1%) was the most common isolate followed by *Aspergillus* spp (28.93%). *Aspergillus fumigatus* was most frequently isolated among *Aspergillus* spp the other common isolate were *Penicillium* spp (7.89%), *Candida albicans* (7.89%), *Curvularia* spp (5.26%), *Cladosporium* spp (2.63%). These observations were comparable

with studies by Gopinath et al, M J Bharathi et al.<sup>9,20</sup> *Aspergillus* spp was the predominant isolate as compared to *Fusarium* spp in the various studies by Samar K Basak et al(59.8%), Jagdish Chander et al (41.18%), Jagdish Chander et al (35%), Javadi et al (34.8%), Shokohi et al (33.3%).<sup>5,16,20,23,10</sup> Gopinath U et al stated that reports from South India, Florida, and Ghana have identified species of *Fusarium* and *Aspergillus* as common corneal pathogens.<sup>22</sup> *Curvularia* spp were foremost among the pigmented fungi. Vajpayee et al noted that cultures were positive in 51.46% cases, although the fungus was demonstrated by KOH in 94.3% of the culture proven cases.<sup>24</sup> In our study yeasts play a minor role because *Candida* infection predominates in colder climates and is preceded by corneal disease, local immunosuppression or systemic diseases like diabetes mellitus<sup>12</sup>.

In the present study, 16 samples remained sterile on culture despite positive direct microscopic examination. The reasons for cultures to be sterile could be patients already using topical steroids or antifungal agents before corneal scrapings were performed and also sometimes insufficient material was obtained for inoculation.

## 6. Conclusion:

Central corneal ulceration leading most often to unioocular blindness continues to be a common ophthalmologic problem. Keratomycosis is the leading type of the corneal ulcers, with a male predominance; agricultural activity and related ocular trauma being the most common predisposing factors. Direct microscopy with CFW is noted to be more sensitive in detecting fungal elements than fungal culture. Early and rapid identification of the pathological organism is the key to ensuring successful medical therapy for corneal infections and is particularly true for fungal keratitis.

## 7. Acknowledgement

I thank Staff and Postgraduate students of Department of Ophthalmology for their kind cooperation during the study. I also would like to thank the department of Pathology for their kind assistance in using the Fluorescent Microscope.

## 8. References

- [1] Agarwal V et al, Current perspectives in infectious Keratitis. *Indian J ophthalmology* 1994; 42:171-191
- [2] World health organization- Visual impairment and blindness, Fact sheet N 282, October 2011.
- [3] Renuka Srinivasan, Reba Kanungo and Goyal JL. Spectrum of Oculomycosis in South India. *Acta Ophthalmologica* 1991;69: 744-749.
- [4] Jagdish Chander, *Oculomycosis Text Book of Medical Mycology*, 3rd Edn, pp 400 – 404. Mehta Publishers A-16 (East) Naraina II, New Delhi - 110028, India.
- [5] Samar K Basak, Sukumar Basak, Ayan Mohanta, Arun Bhowmick – Epidemiological and microbiological diagnosis of Suppurative Keratitis in Gangetic West Bengal, Eastern India. *Indian J Ophthalmol* 2005; 53: 17-22.
- [6] Jayahar Bharathi M, Ramakrishnan R, Samala vasu, Meenakshi R, Palaniappan R- Epidemiological characteristics and laboratory diagnosis of fungal keratitis. *Indian J ophthalmol* 2003; 51:315-21
- [7] Campbell M.C, Joyce L Stewart – “The Medical Mycology hand book”. John Wiley & Sons, New York.1980.436. (173,361).
- [8] McLeod SD; Infectious keratitis in Yanoff M, Duker JS. *Ophthalmology*, 2nd edition, St. Louis Mosby Company, 2004.
- [9] Bharathi MJ, Ramakrishnan R, Meenakshi R, Mittal S, Shivakumar C, Srinivasan - Microbiological diagnosis of infective keratitis comparative evaluation of direct microscopy and culture results. *Br J Ophthalmol* ; 2006; 90: 1271 – 6.
- [10] Tahereh Shokohi, Kiumars Nowroozpoor, Dailami – Fungal Keratitis in patients with corneal ulcer in San, Northern Iran. *Archives of Iranian Medicine* 2006; 9: 222 – 227.
- [11] Weihong Zhang, Huashan Yang, Lili Jiang, Llya Wang- Use of potassium hydroxide, Giemsa and Calcofluor white staining techniques in the Microscopic Evaluation of Corneal scrapings for diagnosis of fungal keratitis. *J International Medical research* 2010; 38: 1961-67
- [12] Sharma S; Silverberg M, Mehta P, Gopinathan U, Naduvilath L.V Prasad eye institute Hyderabad, India – early diagnosis of mycotic keratitis: Predictive value of KOH preparation. *Indian J Ophthalmol* 1998: 46: 31 – 5.
- [13] Thomas PA –Fungal infections of the cornea. *Eye* 2003; 17: 852-62
- [14] Smolin G, Thoft RA, *The Cornea* 3rd edition, Little, Brown and Company, Boston USA, 1994.
- [15] Whitcher JP, Srinivasan M, Upadhyay MP. Prevention of corneal ulceration in the developing world. *Int Ophthalmol clin* 2002; 42(1): 71-77.
- [16] Jagdish Chander, Nidhi Singla, Nalini Agnihotri, Sudesh Kumar Arya, Antariksh Deep - Keratomycosis in and around Chandigarh ; A five year study from a north Indian tertiary care hospital. *Indian J Patho-Microbiol* 2008; 51: 2:304 – 306.
- [17] Sharma S et al – evaluation of corneal scraping smear examination methods in the diagnosis of bacterial and fungal keratitis survey of eight years of laboratory experience. *Cornea* 2002; 21(7):643-7
- [18] Srinivasan M, Christine A, Gonzales, Celine George, Vicky Cevallos, Jeena M, Mascarenhas, Asokan B, John Wilkins, Gilbert Smolin, John P, Whitcher, “Epidemiology and etiological diagnosis of corneal ulceration in Madurai, South India, *Br. J. ophthalmol* 1997:81:965-971.
- [19] Chowdhary A, Singh K. spectrum of fungal keratitis in north India. *Cornea* 2002; 21:161-163.
- [20] Chander J, Chakrabarti A, Shama A, Saini JS, Panigarhi D - Evaluation of calcofluor white staining in the diagnosis of fungal corneal ulcer. *Mycosis* 1993; 36: 243 – 245.
- [21] Savithri Sharma, Silverberg, Paras, Usah Gopinathan, Thomas; Early diagnosis of Mycotic Keratitis: Predictive value of potassium hydroxide preparation *Ind J Ophthalmol*; 1998:46: 31
- [22] Gopinathan U et al- The epidemiological features and laboratory results of fungal keratitis; 10 year review at a referral eye care centre in South India. *Cornea*, 2002; 21 (6): 555-9
- [23] Vajpayee R.B et.al – Laboratory diagnosis of keratomycosis: comparative evaluation of direct microscopy and culture results. *Ann Ophthalmol*, 25 (2): 68-71.